

ASYMPTOTIC PROPERTIES OF INDUCED MAXIMUM LIKELIHOOD  
ESTIMATES OF NONLINEAR. (U) ADVANCED STATISTICAL  
TECHNOLOGIES CORP LAWRENCEVILLE NJ D H JONES OCT 85  
ROB-83-85 N00014-83-C-0627 F/G 12/1


NL

**UNCLASSIFIED**

ROB-03-85 N00014-83-C-0627

F/G 12/1

END



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A163 518

12

ASYMPTOTIC PROPERTIES OF INDUCED MAXIMUM LIKELIHOOD ESTIMATES OF  
NONLINEAR MODELS FOR ITEM RESPONSE VARIABLES:  
THE FINITE-GENERIC-ITEM-POOL CASE

Douglas H. Jones

Technical Report No. ONR-01-85

Advanced Statistical Technologies Corporation  
P.O. Box 6640  
Lawrenceville, New Jersey 08648

October 1985

DTIC  
ELECTE  
JAN 31 1986  
S D

Prepared under contract No. N00014-83-C-0627, NR 150-152  
with the Personnel and Training Research Programs  
Psychological Sciences Division  
Office of Naval Research

Approved for public release: distribution unlimited.

Reproduction in whole or in part is permitted for any purpose  
of the United States Government.

86 1 31 0 2 8

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report No. ONR-01-85	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Asymptotic properties of induced maximum likelihood estimators of nonlinear models for item response variables: the finite-generic-item-pool case		5. TYPE OF REPORT & PERIOD COVERED Technical Report 01 OCT 84 - 31 JUL 85
		6. PERFORMING ORG. REPORT NUMBER ROB-03-85
7. AUTHOR/s  Douglas H. Jones		8. CONTRACT OR GRANT NUMBER(s)  N00014-83-C-0627
9. PERFORMING ORGANIZATION NAME AND ADDRESS Advanced Statistical Technologies Corporation P.O. Box 6640 Lawrenceville, NJ 08648		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61153N, RR4204, RR0420401, NR 150-152
11. CONTROLLING OFFICE NAME AND ADDRESS Personnel and Training Research Programs Office of Naval Research, Code 442PT Arlington, VA 22217		12. REPORT DATE 01 AUGUST 1985
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 32
		15. SECURITY CLASS. (of this report)  Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Item Response Theory, Robustness, Ability, Maximum Likelihood, Influence Curve, Jackknife, Breakdown Point, Estimation.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The progress of modern mental test theory depends very much on the techniques of maximum likelihood estimation, and many popular applications make use of likelihoods induced by logistic item response models. While, in reality, item responses are nonreplicate within a single examinee and the logistic models are only ideal, practitioners make inferences using the asymptotic distribution of the maximum likelihood estimator derived as if item responses were replicate and satisfied their ideal model. This article proposes a sample space acknowledging		

DD FORM 1473

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

these two realities and derives the asymptotic distribution of the induced maximum likelihood estimator.

This article assumes that items, while sampled from an infinite set of items have but a finite domain of alternate response functions: this situation is the case of the finite-generic-item-pool. Later articles will attempt to remove this assumption.

Using the proposed sample space, the article applies the statistical functional approach of von Mises to derive the influence curve of the maximum likelihood estimator; to discuss related robustness properties; and to derive new classes of resistant estimators. This article's general purpose is revealing the value of these methods for uncovering the relative merits of different item response functions. Proofs and mathematical derivations are minimized to increase the assessability of this complex subject.

S N 0102- LF- 014- 6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

14

**Asymptotic properties of induced maximum likelihood estimators  
of non-linear models for item response variables:  
the finite-generic-item-pool case**

Douglas H. Jones  
Advanced Statistical Technologies Corporation

**Abstract**

The progress of modern mental test theory depends very much on the techniques of maximum likelihood estimation, and many popular applications make use of likelihoods induced by logistic item response models. While, in reality, item responses are nonreplicable within a single examinee and the logistic models are only ideal, practitioners make inferences using the asymptotic distribution of the maximum likelihood estimator derived as if item responses were replicated and satisfied their ideal model. This article proposes a sample space acknowledging these two realities and derives the asymptotic distribution of the induced maximum likelihood estimator.

This article assumes that items, while sampled from an infinite set of items, have but a finite domain of alternate response functions: this situation is the case of the finite-generic-item-pool. Later articles will attempt to remove this assumption.

Using the proposed sample space, the article applies the statistical functional approach of von Mises to derive the influence curve of the maximum likelihood estimator; to discuss related robustness properties; and to derive new classes of resistant estimators. This article's general purpose is revealing the value of these methods for uncovering the relative merits of different item response functions. Proofs and mathematical derivations are minimized to increase the accessibility of this complex subject.

Approved for public release; distribution unlimited.

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

## 1. INTRODUCTION

While maximum likelihood procedures are popular in item response theory (IRT), (Lord, 1980), their insensitivity to departures from assumptions is serious enough to warrant cautious use and further study (Wainer-Wright, 1980; Jones, 1982). The purpose of this article is to explore the behavior of the procedures when the model is not true.

To apply some of the concepts of robustness theory, we found that some of the more important concepts required reformulating the maximum likelihood procedures. In particular the study of the robustness of the maximum likelihood estimator (MLE) requires viewing it as a function of the empirical probability distribution function (PDF). The original formulation of item response theory, as a regression problem, does not allow the summarization of the data in terms of an empirical PDF. In §2, we recast the structure of the problem so that the data can be replaced by an empirical PDF and we reformulate the MLE as a function of it.

In §3, we derive the asymptotic distribution of the MLE when the true PDF is not generated by the assumed model. These results are basic to understanding the sensitivity of the MLE to departures from assumptions. They make heavy use of von Mises's approach to statistical functions (Fillippova, 1982).

In §4, we apply the asymptotic formulas derived in §3 to three popular item response models. A measure of goodness-of-fit, compatible with the MLE, is employed to play the role of the mean squared error (McCullagh-Nelder, 1983; Pregibon, 1981). These results reveal that certain item response models reverse the scale of ability.

In §5, we formulate the basic robustness criteria associated with Hampel's influence curve (IC) (Hampel, 1974; Welsch-Krasker, 1982). We derive a relation between the IC and the maximum bias of the MLE as the true PDF is varied within an  $\epsilon$ -contamination neighborhood of the modeled PDF (Huber, 1981). We also derive the breakdown point (Huber, 1981) of the MLE for certain types of departures from the assumptions. The analysis of these criteria shows how the notion of robustness in IRT is fundamentally different from linear and logistic regression problems.

## 2. GENERAL NOTATION AND STRUCTURE

The basic formulation of IRT based on maximum likelihood is:  $u=1$  (correct) or  $u=0$  (incorrect) is observed for each item  $i$  with likelihood, given a real latent parameter  $\theta$ , equal to

$$h_i(u; \theta) = P_i(\theta)^u [1 - P_i(\theta)]^{1-u}$$

and with  $P_i(\theta)$  the  $i^{\text{th}}$  item response model. The total likelihood based on data  $u_1, u_2, \dots, u_n$  and models  $P_1, P_2, \dots, P_n$  is:

$$L(\theta; u_1, \dots, u_n, P_1, \dots, P_n) = \prod_{i=1}^n h_i(u_i; \theta).$$

For robustness studies, we need to allow for the possibility that the item response models are inaccurate. Thus, we assume that  $E(u_i) \neq P_i(\theta)$ . But we retain the assumption of local independence and call  $P_i(\theta)$  an operational model.

To accommodate items with different difficulties and discriminating powers, and simultaneously, apply standard asymptotic theory, we formulate the sample space as:

$$(\text{Sample Space}) \quad S = \{(u, x) : u=0,1; x \in X\}$$

$$X = \text{finite set indexing items.}$$

An observation on  $S$  is denoted by  $s$  or  $t$ , etc., and is generated by administering a randomly chosen item,  $x$ , to obtain a response,  $u$ .



An arbitrary probability distribution function (PDF) on  $S$  is denoted by  $\eta$ . A probability distribution over  $X$  is denoted by  $p$ . The conditional probability distribution of  $u$  given  $x$  is denoted by  $f(u;x)$ . For arbitrary  $\eta$ , there is a  $p$  and  $f(u;x)$  such that:

$$\eta(s) = f(u;x) p(x), \quad s = (u,x).$$

Because  $u$  is binary;  $f(u;x)$  is Bernoulli with some probability of success,  $\Pi^*(x)$  satisfying:

$$f(u;x) = \Pi^*(x)^u [1-\Pi^*(x)]^{1-u}.$$

The empirical PDF defined for a sample  $s_1, s_2, \dots, s_n$  is defined by denoting  $\delta_s$  to be a point mass at  $s$  and

$$\hat{\eta}_n(t) = n^{-1} \sum_{i=1}^n \delta_{s_i}(t).$$

It is a PDF on  $S$ . The distance between two PDF's  $\xi$  and  $\eta$  is defined as  $|\xi - \eta| = \max_{s \in S} |\xi(s) - \eta(s)|$ .

A parametric family of PDF's on  $S$  is defined by  $\{\eta_\theta; \theta \text{ real}\}$ . Values of  $\eta_\theta$  are denoted by  $\eta(s; \theta)$ . A special type of a parametric family is generated by a set of operational models:

$$\text{Operational Models: } \{\Pi(\theta; x) : x \in X; \theta \text{ real}\}$$

$$\text{Parametric Family: } f(u; \theta, x) = \Pi(\theta; x)^u \{1 - \Pi(\theta; x)\}^{1-u}$$

$$\eta(s; \theta) = f(u; \theta, x) p(x).$$

The traditional structure of IRT is related as follows: the  $i^{\text{th}}$  observation is  $u_i$  with model  $P_i(\theta)$ . Let  $x_i$  be the index value of the  $i^{\text{th}}$  chosen item where  $\Pi(\theta; x_i) = P_i(\theta)$ . Let  $s_i = (u_i, x_i)$ , so that  $\eta(s_i; \theta) = f(u_i; \theta, x_i) p(x_i) = h_i(u_i; \theta) p(x_i)$ .

The likelihood based on the sample  $s_1, s_2, \dots, s_n$  is:

$$L(\theta; s_1, s_2, \dots, s_n) = \prod_{i=1}^n \eta(s_i; \theta).$$

If  $\{p(x_i): i=1, \dots, n\}$  contains no information about  $\theta$ , MLE's based on the two likelihoods are identical.

The log-derivative of the parametric PDF is denoted by  $\ell(s; \theta) = (d/d\theta) \log \eta(s; \theta)$ . If it exists, a solution of the implicit equation

$$(\text{Normal Equation}) \quad 0 = \sum_{i=1}^n \ell(s_i; \theta)$$

is denoted by  $\hat{\theta}_n$  and is called the MLE. This equation simplifies with operational models as follows: The logit of an operational model  $\Pi(\theta; x)$  and its derivative is

$$g(\theta; x) = \log \Pi(\theta; x) / [1 - \Pi(\theta; x)]$$

$$g'(\theta; x) = v(\theta; x)^{-1} \Pi'(\theta; x), \text{ where}$$

$$v(\theta; x) = \Pi(\theta; x) [1 - \Pi(\theta; x)].$$

Using the definition of  $\eta_\theta$ , we have:

$$\ell(s; \theta) = g'(\theta; x) [u - \Pi(\theta; x)]$$

and the normal equation becomes

$$0 = \sum_{i=1}^n g'(\theta; x_i) [u_i - \Pi(\theta; x_i)] = \sum_{i=1}^n v(\theta; x_i)^{-1} [u_i - \Pi(\theta; x_i)] \Pi'(\theta; x_i)$$

The Fisher information of the parametric PDF,  $\eta_\theta$ , is

$$I(\theta) = -\sum \ell'(s; \theta) \eta(s; \theta) = \sum \ell(s; \theta)^2 \eta(s; \theta)$$

where the sum is over all  $s$  in  $S$ . This information identity follows from the total differential of  $0 = \sum \ell(s; \theta) \eta(s; \theta)$ ; using  $\eta'(s; \theta) = \eta(s; \theta)(d/d\theta) \log \eta(s; \theta) = \eta(s; \theta) \ell(s; \theta)$  we have,

$$\begin{aligned} 0 &= \sum \ell'(s; \theta) \eta(s; \theta) + \sum \ell(s; \theta) \eta'(s; \theta) \\ &= \sum \ell'(s; \theta) \eta(s; \theta) + \sum \ell(s; \theta)^2 \eta(s; \theta). \end{aligned}$$

Note for computational purposes:  $I(\theta) = \sum g'(\theta; x)^2 v(\theta; x) p(x)$ .

**Example.** The one-parameter logistic (1PL) and two-parameter logistic (2PL)

item response models are characterized by their logits:

$g(\theta; x) = a(x)[\theta - b(x)]$  where  $a(x) > 0$ ,  $-\infty < b(x) < \infty$  are the

discrimination and difficulty parameters for item  $x$ . Hence,

$\ell(s; \theta) = a(x)[u - \Pi(\theta; x)]$  and  $0 = \sum_{i=1}^n a(x_i)[u_i - \Pi(\theta_i; x_i)]$  is the normal

equation. The Fisher information is  $I(\theta) = \sum a(x)^2 v(\theta; x)p(x)$ , sum over all  $X$ .

We wish to generalize the normal equation in two ways: first, we want to show the explicit relation between  $\hat{\theta}_n$  and  $\hat{\eta}_n$ ; second, we wish to consider estimators that are more general than MLE's.

We rewrite the normal equation using the empirical PDF as

$$0 = \sum \ell(s; \theta) \hat{\eta}_n(s)$$

where it will be understood that the sum is always over  $S$ . We see that the MLE depends explicitly on the empirical PDF, we denote this dependence by

$$\hat{\theta}_n = \theta(\hat{\eta}_n).$$

If the empirical PDF is replaced by an arbitrary PDF, the normal equation defines a general functional relationship,  $\theta(\eta)$ , between  $\theta$  and  $\eta$ : we call  $\theta(\eta)$  a statistical functional.

We define M-type estimators generated by a score function  $\psi(s; \theta)$  by the equation

$$0 = n^{-1} \sum_{i=1}^n \psi(s_i; \theta) = \sum \psi(s; \theta) \hat{\eta}_n(s).$$

We see that  $\psi(s; \theta) = \ell(s; \theta)$  generates the MLE. We add this generality because our methods of proof in the next section are really about M-type estimators with the MLE results following as a special case. Note that the notion of a statistical function applies to M-type estimators also. More definitions that we need follow.

We define

$$m(\theta, \eta) = \sum \psi(s; \theta) \eta(s)$$

for an arbitrary PDF and score function. The derivative of  $m(\theta, \eta)$  with respect to  $\theta$  is  $m'(\theta, \eta)$ . Note that  $m'(\theta, \eta_0)$  means  $m'(\theta, \eta)$  evaluated with  $\eta = \eta_0$ . The normal equation is  $0 = m(\theta, \hat{\eta}_n)$  and Fisher's information is  $I(\theta) = -m'(\theta, \eta_0)$  with  $\psi = \ell$ . The Newton-Rapheson algorithm for solving the normal equation is

$$\theta^{t+1} = \theta^t + m(\theta^t, \hat{\eta}_n) / -m'(\theta^t, \hat{\eta}_n);$$

if  $\psi = \ell$  the Fisher scoring algorithm is

$$\theta^{t+1} = \theta^t + m(\theta^t, \hat{\eta}_n) / I(\theta^t).$$

Let  $\psi(s; \theta)$  be a given score function and let  $\theta_0$  denote the value of  $\theta$  that solves the equation  $0 = m(\theta, \eta)$ , corresponding to this score function. If the PDF  $\eta$  is a member of some parametric family and satisfies  $\eta = \eta_{\theta_1}$  for a given fixed parameter value  $\theta_1$  and if  $\theta_0 = \theta_1$ , then we say that the score function is unbiased.

If  $\psi$  is an unbiased score function, then  $0 = m(\theta, \eta_0)$  for all  $\theta$ . This fact leads to an identity that is analogous to the Fisher information identity presented previously and is proven in exactly the same way. The identity is:

$$-m'(\theta, \eta_0) = \sum \psi(s; \theta) \ell(s; \theta) \eta(s; \theta).$$

If one replaces  $m'(\theta, \hat{\eta}_n)$  by its expectation under  $\eta_0$  in the Newton-Rapheson algorithm, one obtains an algorithm that is analogous to Fisher scoring. If  $\psi$  is an unbiased score function, then one may use the above identity for  $-m'(\theta, \eta_0)$  to avoid evaluating the derivative of  $\psi$ .

An important subclass of unbiased score functions are generated by an arbitrary weight function  $w(\theta; x)$  where

$$\psi(s; \theta) = w(\theta; x) [u - \Pi(\theta; x)] \Pi'(\theta; x).$$

If we choose

$$w(\theta; x) = v(\theta; x)^{-1}$$

then  $\psi(s; \theta) = \ell(s; \theta)$  and we are back to the MLE. Other choices of the weight function lead to resistant estimators. For example, Jones (1982) suggests  $w(\theta; x) = v(\theta; x)^{h-1}$  with  $h \geq 0$ , a tuning constant. We can stay in the class of exponential families with arbitrary response variable  $u$ , as long as  $\Pi(\theta; x) = E(u | \theta, x)$  and  $w(\theta; x) = \text{var}(u | \theta, x)^{-1}$  (see Jennrick and Moore, 1975).

Jones' resistant estimator could be generated for these families also by letting  $w(\theta; x) = \text{var}(u | \theta, x)^{h-1}$ . We obtain a more general class of estimators by allowing the weight functions to depend on the response:  $\psi(s; \theta) = w(\theta; s) [u - \Pi(\theta; x)] \Pi'(\theta; x)$ . Krasker and Welsch (1982) consider these estimators for the general linear model. Stefanski, Carroll, and Ruppert (1984) consider these estimators for the logistic model.

An algorithm based on Gauss-Newton's algorithm for solving normal equations with score  $\psi(s; \theta) = w(\theta; s) [u - \Pi(\theta; x)] \Pi'(\theta; x)$  is as follows: (see Holland and Welsch, 1977) define  $d_i^t = \Pi'(\theta; x_i)$  and  $w_i = w(\theta_i; s_i)$  then

$$\theta^{t+1} = \theta^t + [\sum d_i^t w_i d_i^t]^{-1} \sum d_i^t w_i [u_i - \Pi(\theta^t; x_i)].$$

This algorithm is iterative reweighted least squares: at convergence  $\hat{\theta}_n = \theta^\infty$ , so if we define the pseudo-observation  $z_i = d_i \hat{\theta}_n + [u_i - \Pi(\hat{\theta}_n; x_i)]$  then

$$\hat{\theta}_n = [\sum d_i w_i d_i]^{-1} \sum d_i w_i z_i.$$

See Pregibon (1981) and McCullagh and Nelder (1983) for a similar algorithm based on the exponential family with linear predictors. Note that this algorithm is also identical to Fisher scoring.

### 3. GENERAL ASYMPTOTIC THEORY OF M-TYPE ESTIMATORS

We present consistency and asymptotic normality (AN) results in this section. In the first part we confine attention to the main results and in the second part we supply the proofs. Readers may skip the proofs and move on to the next section. In the main results we discuss conditions for consistency and AN. We also characterize an approximation to the M-type estimator that is important for AN results and for the robustness results in §5.

#### 3.1 Main Results: Consistency

Suppose the sample  $s_1, s_2, \dots, s_n$  is IID with PDF  $\eta$ . The empirical PDF  $\hat{\eta}_n$  satisfies  $|\hat{\eta}_n - \eta| \rightarrow 0$  wp 1 as  $n \rightarrow \infty$ . This fact gives us the obvious candidate for the limit of an M-type estimator,  $\theta_0$  which solves  $0 = m(\theta_0, \eta)$ ; when does  $\hat{\theta}_n \rightarrow \theta_0$ ? It is possible that the equation  $0 = m(\theta, \hat{\eta}_n)$  yielding  $\hat{\theta}_n$  has more than one solution, in which case a consistency result may be about only one of the possible sequences of M-type estimators. Also, it is possible that the equation  $0 = m(\theta, \eta)$  does not have a local solution, in which case a consistency result would not be useful. Some known results follow.

Huber (1964): Let  $\theta_0$  be the unique solution. If  $\psi(s; \theta)$  is monotone in  $\theta$  for each  $s \in S$  then every sequence  $\hat{\theta}_n \rightarrow \theta_0$  wp1.

Boos (1977): Let  $\theta_0$  be an isolated solution. If  $\psi(s, \theta)$  is continuous in  $\theta$  for each  $s \in S$  then there exists a sequence  $\hat{\theta}_n \rightarrow \theta_0$  wp1.

Huber (1967, 1980): Let  $\theta_0$  be a unique solution. Let  $|m(\theta, \eta)|$  be bounded from zero as  $|\theta| \rightarrow \infty$ . If  $\psi(s; \theta)$  is continuous in  $\theta$  for each  $s \in S$ , then every sequence  $\hat{\theta}_n \rightarrow \theta_0$  wp1.

The various conditions for consistency will be satisfied when we impose stricter conditions for AN.

**Example.** For the 2PL model  $\Pi(\theta; x)$  is strictly monotone increasing and hence

$\psi(s; \theta) = a(x)[u - \Pi(\theta; x)]$  is monotone, the solution  $\theta_0$  to  $0 = m(\theta_0, \eta) = \int a(x)[\Pi^*(x) - \Pi(\theta_0; x)]p(x)$  is unique with  $\Pi^*(x)$  arbitrary. Thus for  $\eta$  arbitrary, Huber (1964) applies. If  $\eta = \eta_{\theta_1}$  for some fixed  $\theta_1$ , then  $\theta_0 = \theta_1$  is the unique solution.

### 3.2 Main Results: Asymptotic Normality

The first order asymptotic properties of M-type estimators are characterized by the influence curve. The influence curve (IC) is defined as: let  $s \in S$  and  $\delta_s$  a point mass at  $s$ , for  $\epsilon > 0$

$$IC(s, \eta, \psi) = \lim_{\epsilon \rightarrow 0} \frac{\theta(\eta + \epsilon(\delta_s - \eta)) - \theta(\eta)}{\epsilon}.$$

Denoting  $\theta(\delta_s; \epsilon) = \theta(\eta + \epsilon(\delta_s - \eta))$ , we see that the influence curve is an ordinary derivative of  $\theta(\delta_s; \epsilon)$  evaluated at 0:  $IC(s, \eta, \psi) = (d\theta(\delta_s; \epsilon)/d\epsilon)_0$ . For M-type estimators, it will be proved later that for  $\xi$  an arbitrary PDF,  $\int IC(s, \eta, \psi)\xi(s) = (d\theta(\xi; \epsilon)/d\epsilon)_0$ . This latter characterization allows us in §5 to make an important connection between the bias and the influence curve of MLE's. Also for M-type estimators the influence curve is:

$$(**) \quad IC(s, \eta, \psi) = \psi(s; \theta_0) / -m'(\theta_0, \eta)$$

where throughout the remainder of this section  $\theta_0$  satisfies  $0 = m(\theta_0, \eta)$  and  $m'(\theta_0, \eta) < 0$ .

**Example.** For the 2PL and  $\eta$  arbitrary

$$IC(s, \eta, \psi) = a(x)[u - \Pi(\theta_0; x)] / \int a(x)^2 v(\theta_0; x)p(x).$$

Normally the denominator would depend explicitly on  $\Pi^*(x)$  but does not, since  $g''(\theta; x) = 0$ . However, it does depend on  $\Pi^*(x)$  through  $\theta_0$  which solves  $0 = \int a(x)[\Pi^*(x) - \Pi(x; \theta_0)]p(x)$ .

The primary application we make of the influence curve in this section is to get a leading term approximation:

$$(*) \quad \hat{\theta}_n - \theta_0 = n^{-1} \sum_{i=1}^n IC(s_i, \eta, \psi) + R_n$$

where  $R_n$  is the remainder term. We show below  $n^{\frac{1}{2}} R_n \rightarrow 0$  in probability, thus the behavior of  $n^{\frac{1}{2}}(\hat{\theta}_n - \theta_0)$  is deduced from the approximation and the Lindeberg-Levy central limit theorem. The sufficient conditions for AN of M-type estimators are listed under (C). They are implied by conditions (D) when the M-type estimator is an MLE:

(C) There exists an open interval  $\Omega_0$  and a constant  $c > 0$  such that for all  $\theta$  in  $\Omega_0$

C-1:  $\psi(s; \theta)$ ;  $\psi'(s; \theta)$ ;  $\psi''(s; \theta)$  exist for all  $s \in S$ , with the first two continuous in  $\theta$ ;

C-2:  $m'(\theta_\epsilon, \eta_\epsilon) < 0$  for all  $0 \leq \epsilon \leq 1$  and  $\xi$ :  $|\eta - \xi| \leq \epsilon$  where  $\theta_\epsilon$  solves  $0 = m(\theta, \eta_\epsilon)$  and  $\eta_\epsilon = \eta + \epsilon(\xi - \eta)$ .

(D) Define  $\Omega_0 = \{\theta: v(\theta; x) > 0 \text{ for all } x \in X\}$  then suppose  $\Omega_0$  is not empty and for all  $\theta \in \Omega_0$

D-1:  $\Pi'(\theta; x)$ ;  $\Pi''(\theta; x)$ ;  $\Pi'''(\theta; x)$  exist for all  $x$ , with the first two continuous in  $\theta$ ;

D-2:  $\sum g''(\theta; x) [\Pi^*(x) - \Pi(\theta; x)] p(x) < \sum g'(\theta; x)^2 v(\theta; x) p(x)$  for  $\Pi^*(x)$  in an open interval for each  $x$ .

**Theorem 1.** Assume (C) then  $n^{\frac{1}{2}}(\hat{\theta}_n - \theta_0)$  is AN mean 0 and variance:

$$\sigma_0^2 = \sum IC(s, \eta, \psi)^2 \eta(s).$$

**Corollary.** Assume (C-2) and  $\eta = \eta_{\theta_1}$ ,  $\theta_1$  fixed. If  $\psi$  is unbiased, i.e.  $\theta_0 = \theta_1$ ,

$n^{\frac{1}{2}}[\hat{\theta}_n - \theta_0]$  is AN with mean 0 and variance:

$$\sigma_0^2 = \sum \psi(s; \theta_0)^2 \eta(s; \theta_0) / \left[ \sum \psi(s; \theta_0) \ell(s; \theta_0) \eta(s; \theta_0) \right]^2 \text{ where } \ell(s; \theta) =$$

$(\partial / \partial \theta) \log \eta(s; \theta)$ . Hence an unbiased M-type estimator is efficient

if and only if  $\psi$  is proportional to  $\ell$ , or in other words the MLE is optimal among M-type estimators with unbiased score functions.



### 3.3 Proof of Asymptotic Normality

Define for any  $\xi$  PDF,

$$\eta_\epsilon = \eta + \epsilon(\xi - \eta) \quad \text{and}$$

$$\theta_\epsilon = \theta(\eta_\epsilon).$$

A second order Taylor series expansion about 0 is

$$(***) \quad \theta_\epsilon = \theta_0 + (d\theta/d\epsilon)_0 \epsilon + \frac{1}{2} (d^2\theta/d\epsilon^2)_0 a^2,$$

$0 < a < \epsilon$ . Under conditions (C) we show that this expansion is valid with  $\epsilon=1$ .

Using the expansion with  $\xi = \hat{\eta}_n$ , we obtain the approximation (\*), where  $R_n$  is expressed in terms of the second derivative of  $\theta_\epsilon$ . To show that the IC has the form (\*\*) and that  $n^{\frac{1}{2}} R_n \rightarrow 0$  in probability we derive the first two derivatives as follows.

Define  $M(\theta, \epsilon) = m(\theta, \eta_\epsilon)$ . Because  $\theta_\epsilon$  satisfies  $M(\theta_\epsilon, \epsilon) = 0$  for all  $0 \leq \epsilon \leq 1$ , the first and second total differentials with respect to  $\epsilon$  are identically zero and yield two simultaneous equations involving  $d^j\theta/d\epsilon^j$   $j=1,2$ :

$$(1) \quad (\partial M / \partial \theta)_{\theta_\epsilon} d\theta/d\epsilon + \partial M / \partial \epsilon = 0$$

$$(2) \quad (\partial M / \partial \theta)_{\theta_\epsilon} d^2\theta/d\epsilon^2 + [(\partial^2 M / \partial \theta^2)_{\theta_\epsilon} (d\theta/d\epsilon) + (\partial^2 M / \partial \theta \partial \epsilon)_{\theta_\epsilon}] d\theta/d\epsilon + (\partial^2 M / \partial \epsilon \partial \theta)_{\theta_\epsilon} d\theta/d\epsilon + \partial^2 M / \partial \epsilon^2 = 0.$$

Solving equations (1) and (2) for  $d^j\theta/d\epsilon^j$ ,  $j=1,2$ , and using  $\partial M^2 / \partial \epsilon^2 = 0$ , we have from equation (1):

$$(1') \quad \frac{d\theta}{d\epsilon} = \left( \frac{\partial M}{\partial \epsilon} \right) / - \left( \frac{\partial M}{\partial \theta} \right)_{\theta_\epsilon}$$

and from equations (1') and (2):

$$(2') \quad \frac{d^2\theta}{d\epsilon^2} = \left( \frac{\partial^2 M}{\partial \theta^2} \right)_{\theta_\epsilon} \left( \frac{\partial M}{\partial \epsilon} \right)^2 / - \left( \frac{\partial M}{\partial \theta} \right)_{\theta_\epsilon}^3 + 2 \left( \frac{\partial^2 M}{\partial \epsilon \partial \theta} \right)_{\theta_\epsilon} \left( \frac{\partial M}{\partial \epsilon} \right) / \left( \frac{\partial M}{\partial \theta} \right)_{\theta_\epsilon}^2$$

Now we obtain expression (\*\*) for the IC: Let  $\theta_0 = \theta(\eta)$ , we have

$$\begin{aligned} \partial M / \partial \epsilon &= (\partial / \partial \epsilon) m(\theta_0, \eta_\epsilon) = (\partial / \partial \epsilon) \sum \psi(s; \theta_0) [\eta(s) + \epsilon(\xi(s) - \eta(s))] \\ &= \sum \psi(s; \theta_0) [\xi(s) - \eta(s)] = \sum \psi(s; \theta_0) \xi(s). \end{aligned}$$

Substituting  $(\partial M / \partial \theta)_{\theta_0} = m'(\theta_0, \eta)$

and the last expression into (1') we have for any PDF  $\xi$ ,  $(d\theta/d\epsilon)_0 = \sum \psi(s; \theta_0) \xi(s) / -m'(\theta_0, \eta)$ . Thus for  $\xi = \delta_s$ , we have  $IC(s, \eta, \psi) = \psi(s; \theta_0) / -m'(\theta_0, \eta)$ .

With  $\xi = \hat{\eta}_n$ , the empirical PDF, the MLE  $\hat{\theta}_n = \theta(\eta_1)$  where  $\eta_1$  is  $\eta_\epsilon$  with  $\epsilon=1$ . Using the expansion (\*\*\*) with  $\epsilon=1$  we have

$$\hat{\theta}_n - \theta_0 = n^{-1} \sum_{i=1}^n IC(s_i, \eta, \psi) + R_n$$

where

$$R_n = \frac{1}{2} \left( \frac{d^2 \theta}{d\epsilon^2} \right)_{a^*}, \text{ for some } 0 < a^* < 1.$$

Now we state the conditions for the expansion (\*\*\*) and hence the expression for  $R_n$  to be valid: (see Serfling, 1980, pp. 43, 215).

- (A) Apostle (1957, pg. 96)  $(d\theta/d\epsilon)^+$ , the righthand derivative, and  $d^2\theta/d\epsilon^2$  exist everywhere in the open interval  $(0,1)$ ; with the first continuous in the half-closed interval  $[0,1)$ .

By expression (1') and (2') for  $d\theta/d\epsilon$  and  $d^2\theta/d\epsilon^2$  we have formulated conditions (B) that satisfy conditions (A):

- (B) There exists an open interval  $\Omega_0$  such that for all  $\theta$  in  $\Omega_0$

B-1:  $m(\theta, \eta)$ ,  $m'(\theta, \eta)$ ,  $m''(\theta, \eta)$  exist for all  $\eta$ ;

B-2: there exists a constant  $c$ , such that for all  $\xi$ :  $|\xi - \eta| \leq c$ ,

$m'(\theta_\epsilon, \eta_\epsilon) < 0$  for all  $0 \leq \epsilon \leq 1$ .

To further obtain  $n^{1/2} R_n \rightarrow 0$ , we need to examine the terms in expression (2') and place appropriate conditions on the score function,  $\psi$ . The four terms are:

$$(\partial M / \partial \theta)_{\theta_\epsilon} = m'(\theta_\epsilon, \eta_\epsilon)$$

$$(\partial^2 M / \partial \theta^2)_{\theta_\epsilon} = m''(\theta_\epsilon, \eta)$$

$$(\partial M / \partial \epsilon) = m(\theta_\epsilon, \xi - \eta)$$

$$(\partial M / \partial \epsilon \partial \theta)_{\theta_\epsilon} = m'(\theta_\epsilon, \xi - \eta).$$

These terms apply to  $R_n$  with  $\xi = \hat{\eta}_n$ .

We see that the behavior of  $R_n$  depends directly on that of  $\hat{\eta}_n$  and in particular the differences  $\hat{\eta}_n(s) - \eta(s)$ ,  $s \in S$ . We have given condition (C) at the beginning of this section to keep  $m'(\theta_\epsilon, \eta_\epsilon)$  properly away from zero so as to keep  $R_n$  from exploding and to infer its behavior from that of  $\hat{\eta}_n$ . The following two lemmas imply that  $n^{\frac{1}{2}} R_n \rightarrow 0$  wpl.

**Lemma A.** Assume conditions (C). Put  $\xi = \hat{\eta}_n$  in (2'). There exists constants  $a$  and  $n_0$  such that  $|d^2\theta/d\epsilon^2| \leq a|\hat{\eta}_n - \eta|^2$  for all  $0 \leq \epsilon \leq 1$  and  $n \geq n_0$  wpl.

**Lemma B.** Assume  $s_1, s_2, \dots, s_n$  are IID with PDF  $\eta$ . Then as  $n \rightarrow \infty$  the following holds:

- a)  $\hat{\eta}_n(s) \rightarrow \eta(s)$  for all  $s \in S$  wpl;
- b)  $\{n^{\frac{1}{2}}[\hat{\eta}_n(s) - \eta(s)]: s \in S\}$  converges in law to a Gaussian process with mean 0 and covariance function:

$$\text{COV}(s, t) = \begin{cases} \eta(s)[1-\eta(s)] & s=t \\ -\eta(s)\eta(t) & s \neq t \end{cases};$$

- c)  $|\hat{\eta}_n - \eta| \rightarrow 0$  wpl;
- d)  $n^{\frac{1}{2}}|\hat{\eta}_n - \eta|$  converges in law and in probability.

These two lemmas imply that  $n^{\frac{1}{2}} R_n \rightarrow 0$  in probability and hence  $n^{\frac{1}{2}}(\hat{\theta}_n - \theta_0)$  is AN since lemma A implies  $n^{\frac{1}{2}}|R_n| \leq a n^{\frac{1}{2}}|\eta_n - \eta| \cdot |\eta_n - \eta|$  and lemma B implies that  $|\eta_n - \eta| \rightarrow 0$  while  $n^{\frac{1}{2}}|\hat{\eta}_n - \eta|$  remains bounded in probability.

#### 4. SPECIFIC ASYMPTOTIC BEHAVIOR OF MLE's

Throughout this section  $\eta_0$  will denote a true PDF and  $\theta_0$  the solution to  $0 = m(\theta, \eta_0)$ . The asymptotic behavior of the MLE is obtained from the previous results of M-type estimators with score function  $\ell(s; \theta)$ ,  $s \in S$ . We discuss these

aspects: goodness-of-fit, scale reproduction, and Fisher variance.

Most practical operational models satisfy condition (D-1). Thus, we can say that for situations of interest, MLE's are consistent,  $\hat{\theta}_n \rightarrow \theta_0$  as  $n \rightarrow \infty$  and is asymptotically normal even when  $\eta_0$  is not a member of the parametric family  $\{\eta_\theta: \theta \text{ real}\}$  generated by the set of operational models but satisfies the mild regularity condition (D-2). We will employ this result with the 1PL, 2PL, and 3PL item response models in the example at the end of this section.

Let  $\eta_0$  denote a member of a certain parametric family and consider the MLE associated with the family. If for some  $\theta_1$ ,  $\eta_0 = \eta_{\theta_1}$ , then the MLE is asymptotically unbiased, meaning  $\theta_0 = \theta_1$ . Let  $\xi_{\theta_1}$  denote a member of a different parametric family. If  $\eta_0 = \xi_{\theta_1}$  then the MLE is asymptotically biased meaning  $\theta_0 \neq \theta_1$ . If  $\eta_0$  is not a member of any parametric family then the notion of unbiasedness has no meaning.

Even if  $\eta_0$  is a member of some parametric family not identical to  $\{\eta_\theta\}$ , the bias does hold much information about how good the MLE may be. This is because the parametrization of the family containing  $\eta_0$  is as good as arbitrary when it is not exactly the one generating the MLE. This leads us to propose a different notion of accuracy, possibly supplying the information we usually obtain with measurements of bias.

The information supplied by the bias is obtained from comparison of its square to the variance, because the mean square error, a measure of total error, is the sum of the squared bias and variance. When the bias overwhelms the variance, one usually goes looking for another statistical procedure that can control the bias. (What this compares to in IRT is the adoption of more complex item response models).

A measure that seems to decompose into parts due to "bias" and "variance" and does not depend on the arbitrary parametrization of a true family of PDF's is as follows: for two PDF's  $\eta$  and  $\xi$  define

$$K(\eta, \xi) = \int \eta(s) \log \eta(s)/\xi(s).$$

$K(\eta, \xi)$  is nonnegative and equal to 0 when  $\eta = \xi$ ;  $K'(\eta_0, \eta_\theta) = -m(\theta, \eta_0)$ , thus  $K(\eta_0, \eta_\theta)$  is minimized by  $\theta_0$ . A second order Taylor series expansion gives

$$E K(\eta_0, \eta_{\hat{\theta}_n}) \cong K(\eta_0, \eta_{\theta_0}) - \frac{1}{2} m'(\theta_0, \eta_0) n^{-1} \sigma^2,$$

where we have assumed that the MLE has been modified appropriately to have the moments for the approximation and we have used Theorem 1.

Thus  $E K(\eta_0, \eta_{\hat{\theta}_n})$  behaves as a "total error" and  $K(\eta_0, \eta_{\theta_0})$  behaves as "bias squared" when it is compared to the last term on the right-hand side of the above Taylor series. The quantity  $K(\hat{\eta}_n, \eta_{\hat{\theta}_n})$  is proportional to the deviance in generalized linear models (see McCullagh and Nelder, 1983) and serves as a goodness-of-fit statistic. Values of  $E K(\eta_0, \eta_{\hat{\theta}_n})$  and its approximate components are displayed in Table 3 and discussed in the example at the the end of this section.

Information of a different nature than bias, applicable to arbitrarily parametrized families, is obtained by comparing the rank-order of estimated parameters with the rank-order of known abilities. Suppose there is a certain parametric family  $\{\xi_\theta\}$  of PDF's with the property that  $\eta_0 \in \{\xi_\theta\}$  where  $\eta_0$  may be generated by any member of a population of examinees. But the family  $\{\xi_\theta\}$  is too complex, making its calibration unstable with reasonable sample sizes of examinees. Thus, we prefer instead to use a more parsimonious family  $\{\eta_\theta\}$  with the MLE obtaining  $\theta_0$  as a limit when  $\eta_0 = \xi_{\theta_1}$ ,  $\theta_1$  fixed, and  $n \rightarrow \infty$ . Previous discussion implies that  $\theta_0 \neq \theta_1$ , in general; but the bias here is nonsense. What

is useful is a measure of the distortion between  $\theta_0$  and  $\theta_1$  as  $\theta_1$  moves throughout the population. We do not propose a measure but we do think that one should be sensitive to reversals in the " $\theta$ -scale". Table 3 displays reversals for the 1PL and 2PL families and is further discussed in the example at the end of this section.

Turning from bias to variance, we will now consider the predicament of approximating the true asymptotic variance of an MLE when we do not know  $\eta_0$ . A ready approximation to  $\sigma_0^2$  of Theorem 1 in §3 is the reciprocal of Fisher's information:  $I(\theta_0)^{-1}$  (§2). But we know from §3 that it is not valid when  $\eta_0$  is not a member of the parametric family of PDFs that generate the MLE.

In general, the  $\sigma_0^2$  does not majorize  $I(\theta_0)^{-1}$  or viceversa. Thus, it is possible that the reciprocal of Fisher's information can give either a conservative or misleading approximation to the true asymptotic variance. Table 3 displays the true variance along side the Fisher variance to show both the good and the bad; we discuss this further in the example.

**Example.** Listed in Table 1 are the true and modeled response probabilities of five subjects on four ASVAB items. The true probabilities were actually obtained from a very complex item response model which was calibrated on a very large population. The subjects are ranked from lowest to highest going left to right. The modeled response probabilities follow the 1PL, 2PL, or 3PL item response models as indicated; they too were calibrated on a very large population; Table 2 lists the values of the calibrated parameters. Table 3 is a summary of the asymptotic features of the respective MLE's, which we discuss as follows.

First, note the magnitudes of the traditional notion of bias by taking  $|\theta_1 - \theta_0|$  differences from columns (1) and (2). These values could easily change

upon reparametrization of the true response model; thus they are arbitrary. Thus column (1) should only convey the rank-order of the subjects.

Our notion of "bias squared" is found in column (5),  $K(\eta_0, \theta_0)$ . These values will not change if another parametrization were imposed on the true model. The worst fit is found with subject 5(2PL), referring back to Table 1 we can see that the 2PL model provides poor estimates of all item response probabilities. There are three good values; for example, subject 5(3PL) for which Table 1 shows good estimates of item response probabilities.

Column (3),  $-m'(\theta_0, \eta_0)$ , gives us a feel for the curvature of the likelihood since  $n \cdot m'(\theta_0, \eta_0)$  is an estimate of  $n \cdot m'(\hat{\theta}_n, \hat{\eta}_n)$ , the second derivative of the log-likelihood. We see that the likelihood would tend to be flat for subjects 1,2 (3PL) even though  $K(\eta_0, \theta_0)$  shows close agreement between the estimated and true item response probabilities.

We present the "total error"  $E K(\eta_0, \eta_{\hat{\theta}_n})$ , column (4), for a sample size of  $n=16$ , each item type represented equally. These errors appear to be equal across models and subjects with exception of subject 5 (2PL) as noted before. The components of the total error are in columns (5) and (6) which can tell us the proportion of the total error due to systematic bias:  $(5)/(4)$ . The worst proportion is found with subjects 1, 2 (1PL) meaning that the 1PL is inadequate with these subjects.

We may average the "total error" and "bias squared,"  $E K(\eta_0, \eta_{\hat{\theta}_n})$  and  $K(\eta_0, \eta_{\hat{\theta}_0})$  respectively, over the subjects to get an overall assessment. These averages are for the 1PL, 2PL, and 3PL models respectively:  $(\text{error}, \text{bias}^2) = (.06, .03), (.06, .03), (.04, .01)$ . We see that on average there is at least 25 percent of total error that is systematic bias.

The presence of reversals of the  $\theta_0$ -scale can be detected from column (2). Both the 1PL and 2PL item response models have reversals at the lower abilities. This happens because the 1PL and 2PL calibrations compensate good fit to true response probabilities by distorting the  $\theta_0$ -scale. The Spearman rank correlation between the true ability rank-order and the  $\theta_0$ -scale rank-order of the 1PL, 2PL, and 3PL models are respectively: 0.60, 0.67, 1.00. The numbers may be interpreted as an alternative theoretical goodness-of-fit, since we never would know the true rank-order of ability, there is no practical gain in the measure.

We compare the true variance and the Fisher variance by using columns (7) and (8). For the most part, the Fisher variance yields a conservative assessment of precision; however, it can also be misleading as with subjects 3, 5 (2PL).

**Remarks.** 1) Column (3),  $-m'(\theta_0, \eta_0)$ , can play the role of information. An empirical assessment could be  $-m'(\hat{\theta}_n, \hat{\eta}_n)$ . Also, ratios could play the role of relative efficiency.

2) One should be cautious even if measures of fit, such as  $K(\hat{\eta}_n, \eta_{\hat{\theta}_n})$ , are favorable because as the example shows it is possible to have reversals of the  $\theta_0$ -scale even if the fit is good.

3) We have refrained from making an elaborate comparison of the 1PL, 2PL, and 3PL models based on the data, because one needs to properly account for sampling variability of the calibration process. Such a study is reported in Jones, Wainer and Kaplan (1984).

## 5. SPECIFIC ROBUSTNESS OF THE MLE

Let  $\eta$  denote an arbitrary true PDF,  $\eta_0$  some fixed PDF,  $\{\eta_\theta\}$  a parametric family of PDF's that induces the MLE. Let  $\theta(\eta)$  denote the solution to  $0=m(\theta, \eta)$ .



From Theorem 1 in §3 we have that  $\theta_n \rightarrow \theta(\eta)$  with asymptotic variance  $\sigma^2 = \sigma^2(\eta)$ .

Note that we will not assume that  $\eta$  or  $\eta_0$  belongs to  $\{\eta_0\}$ .

The asymptotic bias of the MLE relative to  $\eta_0$  and  $\{\eta_0\}$  is defined as  $|\theta(\eta) - \theta(\eta_0)|$ . Let  $P_\epsilon$  denote an  $\epsilon$ -neighborhood of  $\eta_0$ , for  $\eta$  belonging to  $P_\epsilon$  we want to quantify the degradation of bias and variance. We say that the robustness of the MLE is measured by the amount of degradation of the maximum bias

$$b(\epsilon) = \sup_{\eta \in P_\epsilon} |\theta(\eta) - \theta(\eta_0)|$$

and the maximum variance

$$v(\epsilon) = \sup_{\eta \in P_\epsilon} \sigma^2(\eta).$$

If  $b(\epsilon)$  were large relative to  $v(\epsilon)$ , then the maximum variance would not be a very important quantifier of robustness. We confine study to  $b(\epsilon)$  in this paper.

There are several important notions for quantifying the robustness of an estimator. Among them are the sensitivities of a parameteric estimator, a fitted value, or a predicted value when one observation is deleted from the sample. These measures are called, respectively, gross error sensitivity (Huber, 1981), change in fit sensitivity and prediction sensitivity (Krasker and Welsch, 1983). The gross error sensitivity is related directly to the maximum bias as shown below. We formulate these quantities and demonstrate their use with the 1PL, 2PL, and 3PL item response models.

Another robustness notion is the sensitivity of the maximum bias as  $\epsilon$  is varied. Certain values of  $\epsilon$  can cause the maximum bias to explode; the smallest such value is called the breakdown point (Huber, 1981). We formulate this quantity and demonstrate its use with the 1PL, 2PL, and 3PL models also.

### 5.1 Sensitivities Based on Deletion

The gross error sensitivity is defined as

$$\gamma^* = \max_s |IC(s, \eta_0, \psi)|.$$

From the leading term approximation of section 3.2, we see that it is proportional to the maximal influence exerted by any one observation on the error of estimation,  $\hat{\theta}_n - \theta(\eta_0)$ . It is related to the maximum bias  $b(\epsilon)$  as follows.

Recall from §3.2 that  $[\theta(\eta_0 + \epsilon(\xi - \eta_0)) - \theta(\eta_0)] \epsilon \rightarrow \sum IC(s, \eta_0, \psi) \xi(s)$  as  $\epsilon \rightarrow 0$ . Let  $P_\epsilon$  be the  $\epsilon$ -contamination neighborhood defined by  $P_\epsilon = \{\eta: \eta = \eta_0 + \epsilon(\xi - \eta_0), \xi \text{ arbitrary PDF}\}$ . Then

$$\sup_{\eta \in P_\epsilon} |\theta(\eta) - \theta(\eta_0)| \cong \epsilon \sup_{\xi} |\sum IC(s, \eta_0, \psi) \xi(s)|.$$

Thus

$$b(\epsilon) \cong \epsilon \gamma^*.$$

So that for small  $\epsilon$ ,  $\gamma^*$  measures the rate of growth of the maximum bias over the  $\epsilon$ -contaminated neighborhood.

For M-type estimators  $\gamma^* = \infty$  is equivalent to a zero breakdown point, meaning that any departure from  $\eta_0$  will cause the maximum bias to explode. Either condition also implies that the estimator is not continuous at  $\eta_0$  when reviewed as a function of  $\eta$  (assuming, of course, a complimentary topology on the set of PDF's). An estimator is qualitatively robust if it is continuous (Huber, 1981), thus an M-type estimator is not robust if  $\gamma^* = \infty$  or the breakdown point is zero.

The gross error sensitivity also measures the maximum change in the estimator caused by deleting one observation. Let  $\hat{\eta}_n(1)$  and  $\hat{\eta}_n(0)$  denote the empirical PDF with and without  $s_i$ . Let  $\hat{\theta}_n(1)$  and  $\hat{\theta}_n(0)$  denote the corresponding MLE's. Then using the direct definition of the influence curve (§3.2) with  $s=s_i$ ,

$\eta = \hat{\eta}_n(0)$  and  $\epsilon = 1/n$  it is easy to show

$$\theta_n(1) - \theta_n(0) \cong n^{-1} IC(s_i, \eta_n(0), \psi).$$

Thus

$$\max_i |\hat{\theta}_n(1) - \hat{\theta}_n(0)| \cong n^{-1} \gamma^*.$$

The change in fit sensitivity concerns the effect of deleting one observation,  $s_i$ , on the estimated logit,  $g(\hat{\theta}_n; x_i)$ . This change in fit is  $g(\hat{\theta}_n(1); x_i) - g(\hat{\theta}_n(0); x_i) \cong g'(\hat{\theta}_n(0); x_i) [\hat{\theta}_n(1) - \hat{\theta}_n(0)]$ . Putting this together with the estimator sensitivity we have

$$g(\hat{\theta}_n(1); x_i) - g(\hat{\theta}_n(0); x_i) \cong n^{-1} g'(\hat{\theta}_n(0); x_i) IC(s_i, \hat{\eta}_n(0), \psi).$$

Thus the shape of  $g'(\theta; x) IC(s, \eta, \psi)$  would indicate robustness as would the size of the change in fit sensitivity:

$$\gamma^{**} = \max_s |g'(\theta; x) IC(s, \eta, \psi)|.$$

Prediction sensitivity concerns the effect of deleting an observation from the sample on the predicted logit of the future item,  $g(\hat{\theta}_n; z)$  where  $z$  is yet to be administered. Let  $\lambda = g'(\theta; z)$ , then by a Taylor series approximation,  $g(\hat{\theta}_n; z) \cong g(\theta; z) + \lambda(\hat{\theta}_n - \theta)$ . Hence the change in prediction is measured by the change in  $\lambda \hat{\theta}_n$ , and  $\lambda IC(s_i, \eta, \psi)$  measures this change due to deleting  $s_i$ . To be meaningful this change must be weighed relative to its standard deviation,  $\lambda [\sum IC(s, \eta, \psi)^2 \eta(s)]^{\frac{1}{2}}$ . Thus the shape of the ratio indicates robustness as would the prediction sensitivity:

$$\gamma = \max_s \frac{|IC(s, \eta, \psi)|}{[\sum IC(s, \eta, \psi)^2 \eta(s)]^{\frac{1}{2}}}.$$

We can simplify this quantity to show the direct dependence on the score function by using the formula for the influence curve:

$$\gamma = \max_s \frac{|\psi(s;\theta)|}{[\sum \psi(s;\theta)^2 \eta(s)]^{\frac{1}{2}}}$$

where  $\theta$  is evaluated at  $\theta(\eta)$ .

Now we study the various sensitivities to get a feel for their implications in IRT using the 1PL, 2PL, and 3PL models as examples. Graphs of these quantities are useful but require specific values for item parameters and do not lead to any more profound conclusions than just analytic circumspection. Graphs are most useful, however, with actual data, providing diagnostic information on the fit of the model. We study only the MLE induced by  $\{\eta_\theta\}$  and do not look at general M-type estimators. We also restrict this study to sensitivities to departures from the parametric model, that is we let  $\eta_0 = \eta_\theta$  for some value of  $\theta$ . Huber (1980) remarks that a better indication of robustness is to allow  $\eta$  to roam around a  $P_\epsilon$  neighborhood of  $\eta_\theta$  while looking at the sensitivities. We do not have the analytical means to do this at this time.

Consider now and for the rest of this section the MLE with operational models  $\{\Pi(\theta;x): x \in X\}$ . With  $\eta_0 = \eta_\theta$ ,  $-m'(\theta; \eta_\theta) = \sum \psi(s;\theta) \ell(s;\theta) \eta(s;\theta)$  and with  $\psi(s;\theta) = \ell(s;\theta) = g'(\theta;x)[u - \Pi(\theta;x)]$ , we have

$$IC(s, \eta, \ell) = \frac{g'(\theta;x)[u - \Pi(\theta;x)]}{\sum g'(\theta;x)^2 v(\theta;x) p(x)}.$$

Define

$$M(\theta;x) = \max\{\Pi(\theta;x), 1 - \Pi(\theta;x)\},$$

the various sensitivities to departures from  $\eta_0 = \eta_\theta$  are

$$\gamma^* = \frac{\max_x g'(\theta;x) M(\theta;x)}{\sum g'(\theta;x)^2 v(\theta;x) p(x)}.$$

$$\gamma^{**} = \frac{\max_x g'(\theta;x)^2 M(\theta;x)}{\sum g'(\theta;x)^2 v(\theta;x) p(x)} \quad \text{and}$$

$$\gamma = \frac{\max_x g'(\theta; x) M(\theta; x)}{[\sum g'(\theta; x)^2 v(\theta; x) p(x)]^{\frac{1}{2}}}.$$

**Example 1.** The 2PL item response models have  $g'(\theta; x) = a(x)$  where  $a(x) > 0$ . If  $a(x) \equiv a_0$ , then the models are called the 1PL item response models.  $\gamma^*$  is finite provided  $\max a(x)$  is finite. If the generic item pool  $X$  is finite then  $\gamma^*$  is always finite. If the generic item pool is not finite, it is possible that  $\sup a(x) = \infty$  but practical reasons would disallow this from happening because an "infinitely discriminating item" is rare.

**Example 2.** The 3PL item response models are defined as  $\Pi(\theta; x) = [1 - c(x)] R(\theta; x) + c(x)$  where  $0 < c(x) < 1$  and  $R(\theta; x)$  is a 2PL model. Define  $v_1(\theta; x) = R(\theta; x)[1 - R(\theta; x)]$ . It can be shown that  $g'(\theta; x) = [1 - c(x)] a(x) v_1(\theta; x) / v(\theta; x)$ .  $\gamma^*$  is finite provided  $\max a(x)$  is finite, the discussion in the previous example applies here too.

**Example 3.** For all the 1PL, 2PL, and 3PL models, because of the behavior of  $g'(\theta; x)$ ,  $\gamma^{**}$  and  $\gamma$  are finite if and only if  $\max a(x)$  is finite.  $\gamma^{**}$  and  $\gamma$ , but not  $\gamma^*$ , are invariant for changes of scale in  $a(x)$  and  $b(x)$ . Presumably  $c(x)$  is scale free as it is a probability of the examinee guessing the correct answer to item  $x$ .

The examples lead to the general conclusion that  $\gamma$ ,  $\gamma^*$ , and  $\gamma^{**}$  are finite if and only if  $\max |g'(\theta; x)|$  is finite. For the 1PL, 2PL, and 3PL models this condition is equivalent to having  $\max a(x)$  finite.

Because the sensitivities change as  $\theta$  changes, their variation over the entire range of practical  $\theta$ -values should be studied to properly assess robustness in IRT. This allows for the fact that the MLE procedure must estimate unique  $\theta$  parameters for different subjects. This is in marked contrast with estimation in logistic regression -- the same estimation procedures as IRT but

the object is to estimate a single  $\theta$  (such as lethal dose 50 or the vector of parameters in one response function). Because the sensitivities must be viewed globally, procedures that are robust for logistic regression may not be directly transferable to IRT.

Consider what happens as  $|\theta| \rightarrow \infty$ . The denominator of  $\gamma^*$  and  $\gamma^{**}$  is Fisher's information; for  $\gamma$ , it is just the square root. For extreme  $\theta$ 's it is reasonable to assume that any finite set of generic items  $X$ , item responses hold little information about  $\theta$ ; thus it is probable that the denominators of the sensitivities approach zero as  $|\theta| \rightarrow \infty$ . Unless the numerators approach zero at the same or faster rate as the denominators, the sensitivities will explode. Applying this idea to each sensitivity, we conclude that  $\gamma^*$  always explodes and for models with  $v(\theta; x) \rightarrow 0$ ,  $\gamma^*$  and  $\gamma^{**}$  both explode. Of the models considered before, the 3PL is the only one having  $v(\theta; x) > 0$  as  $\theta \rightarrow -\infty$ ; thus  $\gamma^*$  and  $\gamma^{**}$  are bounded for the negative extremes of ability.

These results imply that the MLE procedures are not robust because the maximum bias in an  $\epsilon$ -contaminated neighborhood is approximately  $\epsilon\gamma^*$  and  $\gamma^*$  is unbounded as  $|\theta| \rightarrow \infty$ ; thus, the MLE cannot tolerate any contamination at extreme  $\theta$ . The 3PL fares a little better than the 1PL or 2PL as  $\theta \rightarrow -\infty$  since its gross error sensitivity grows a little slower. Thus to achieve full protection one must look outside the class of MLE procedures, which means we have to sacrifice efficiency. (Contrast this with the location problem where the median is the efficient procedure for logistic errors and it is optimal for minimizing the maximum bias; Huber, 1981).

## 5.2 Breakdown Point

The worst possible bias at  $\eta_0$  is defined as  $b(1) = \sup_{\xi} |\theta(\xi) - \theta(\eta_0)|$ , where the supremum is over all arbitrary PDFs,  $\xi$ . Let  $P_{\epsilon}$  be an  $\epsilon$ -neighborhood of  $\eta_0$ .

The breakdown point,  $\epsilon^*$ , is the largest  $\epsilon$  for which  $b(\epsilon)$  is less than the worst value:

$$\epsilon^* = \sup\{\epsilon : b(\epsilon) < b(1)\}.$$

The value of  $\epsilon^*$  depends on the kind of  $P_\epsilon$  chosen; however, it is sometimes adequate to consider just one kind of neighborhood. In IRT,  $b(1) = \infty$ .

We use the following kind of  $\epsilon$ -neighborhood: Let  $0 \leq \Pi \leq 1$  and define  $v = \Pi(1-\Pi)$ . Denote the interval  $D(\theta; x) = [\Pi - \epsilon v^{\frac{1}{2}}, \Pi + \epsilon v^{\frac{1}{2}}]$  when  $\Pi = \Pi(\theta, x)$ ;  $\theta$  is fixed. Denote the subinterval of  $[0, 1]$  by  $D^*(\theta; x) = D(\theta; x) \cap [0, 1]$ . The collection of intervals  $\{D^*(\theta; x)\}$ ,  $x$  fixed, is an  $\epsilon$ -envelope of the item response function  $\Pi(\theta; x)$ . Define  $P_{\epsilon, \theta} = \{\eta : \eta(s) = \Pi^*(x)^u [1 - \Pi^*(x)]^{1-u} p(x); \Pi^*(x) \in D^*(\theta; x)\}$ . It is an  $\epsilon$ -neighborhood "centered" at  $\eta_\theta$ .

Define  $b_+(\epsilon) = \sup_{\xi} [\theta(\xi) - \theta(\eta_0)]$  and  $b_-(\epsilon) = \inf_{\xi} [\theta(\xi) - \theta(\eta_0)]$ . Then  $b(\epsilon) = \max\{b_+(\epsilon), -b_-(\epsilon)\}$ . We consider  $b_+(\epsilon)$  first.

Let  $\eta_0 = \eta_{\theta_0}$ . Define  $\Pi^+(\theta_0; x) = \min\{1, \Pi + \epsilon v^{\frac{1}{2}}\}$  with  $\Pi = \Pi(\theta_0; x)$ . It is clear that  $\Pi^+(\theta_0; x) \in D(\theta_0; x)$  and  $\eta_{\theta_0}^+$ , the corresponding PDF, satisfies  $m(\theta; \eta_{\theta_0}^+) > m(\theta; \eta)$  for all  $\theta$  and all  $\eta \in P_{\epsilon, \theta_0}$ . The maximum "positive" bias satisfies

$$b_+(\epsilon) = \inf\{\theta : m(\theta; \eta_{\theta_0}^+) < 0\} - \theta_0.$$

We have breakdown if  $b_+(\epsilon) = b(1) = \infty$ . To avoid this it is necessary that  $\epsilon$  satisfy  $\lim_{\theta \rightarrow \infty} m(\theta; \eta_{\theta_0}^+) < 0$ . Using the definition of  $m(\theta; \eta)$  we have

$$m(\theta; \eta_{\theta_0}^+) = \sum g'(\theta; x) [\Pi(\theta_0; x) - \Pi(\theta; x)] p(x) + \epsilon \sum g'(\theta; x) v(\theta_0; x)^{\frac{1}{2}} p(x).$$

Letting  $\theta \rightarrow \infty$  and denoting  $g'(\infty; x) = \lim g'(\theta; x)$  we have an equation for the "positive" side breakdown:

$$\epsilon^+ = \frac{\sum g'(\infty; x) [1 - \Pi(\theta_0; x)] p(x)}{\sum g'(\infty; x) v(\theta_0; x)^{\frac{1}{2}} p(x)}.$$

Similarly the "negative" side breakdown is:

$$\epsilon^- = \frac{\sum g'(-\infty; x) \Pi(\theta_0; x) p(x)}{\sum g'(-\infty; x) v(\theta_0; x)^{\frac{1}{2}} p(x)} .$$

And the breakdown:

$$\epsilon^+ = \min(\epsilon^+, \epsilon^-).$$

For a fixed  $\theta_0$ , all MLE procedures for IRT have a breakdown point that is not 0 for  $P_\epsilon$  neighborhoods considered thus far. But as  $|\theta_0| \rightarrow \infty$  the picture changes:  $\epsilon^* = 0$  if either  $\Pi(\theta_0; x) \rightarrow 1$  or 0 for all items  $x$ . Thus the 1PL and 2PL induced MLEs have zero breakdown, meaning they have no tolerance for departures from their models. The 3PL induced MLE has zero breakdown, but for  $\theta \rightarrow \infty$ , the "negative" sided breakdown is not zero, so it could tolerate some departure from its model there.

**Example.** The following displays the "positive" and "negative" breakdown points for the 3PL model with  $a(x) = a_0$  and  $c(x) = c_0$ .

$c_0$	$\epsilon^-$	$\epsilon^+$
.025	.16	0
.05	.23	0
.10	.33	0
.20	.50	0



## REFERENCES

- Apostle, T.M. (1957). Mathematical Analysis. Reading, MA.: Addison-Wesley.
- Boos, D.D. (1977). The differential approach in statistical theory and robust inference. Tallahassee, FL: Florida State University, Ph.D. Dissertation.
- Flippova, A.A. (1962). Mises' theorem of the asymptotic behavior of functionals of empirical distribution functions and its statistical applications. Theor. Prob. App., 7, 24-57.
- Hampel, F.R. (1968). Contributions to the theory of robust estimation. Berkeley, CA: University of California, Ph.D. Dissertation.
- Hampel, F.R. (1974). The influence curve and its role in robust estimation. Journal of the American Statistical Association, 62, 1179-1186.
- Holland, P.W. and Welsch, R.E. (1977). Robust regression using iteratively reweighted least-squares. Communication in Statistics: Theory and Methods, 6, 813-827.
- Huber, P.J. (1964). Robust estimation of a location parameter. Annals of Mathematical Statistics, 35, 73-101.
- Huber, P.J. (1967). The behavior of maximum likelihood estimates with non-standard conditions. In Proceedings of the Fifth Berkeley Symposium on Mathematics Statistics and Probability, Volume 1. Berkeley: University of California Press.
- Huber, P.J. (1981). Robust Statistics. New York: Wiley and Sons.
- Jennrich, R.I. and Moore, R.H. (1975). Maximum likelihood estimation by means of nonlinear least squares. Proceedings of the Statistical Computing Section, 57-65. Washington, DC: American Statistical Association.

- Jones, D.H. (1982). Redescending M-type estimators of latent ability.  
Princeton, NJ: Educational Testing Service, Program Statistics Research  
Technical Report No. 82-30 and Research Report No. 82-15.
- Jones, D.H., Wainer, H., and Kaplan, B. (1984). Estimating ability with three  
item response models when the models are wrong and their parameters are  
inaccurate. Princeton, NJ: Educational Testing Service, Program  
Statistics Research Technical Report No. 84-46 and Research Report No.  
84-26.
- Krasker, W.S. and Welsch, R.E. (1982). Efficient bounded-influence regression  
estimation. Journal of the American Statistical Association, 77, 595-604.
- Lord, F.M. (1980). Application of Item Response Theory to Practical Testing  
Problems. Hillsdale, NJ: Lawrence Erlbaum Associates.
- McCullagh, P. and Nelder, P.A. (1983). Generalized Linear Models. New York:  
Chapman and Hall.
- Pregibon, D. (1981). Logistic regression diagnostics. Annals of Statistics, 9,  
705-724.
- Serfling, R.J. (1980). Approximation Theorems of Mathematical Statistics.  
New York: Wiley and Sons.
- Stefanski, L.A., Carroll, R.J., and Ruppert, D. (1984). Bounding influence and  
leverage in logistic regression. Ithaca, NY: Cornell University,  
Department of Economic and Social Statistics.
- Wainer, H. and Wright, B.D. (1980). Robust estimation of ability in the Rasch  
model. Psychometrika, 45, 373-390.

TABLE 1. ITEM RESPONSE PROBABILITIES

MODEL	ITEM 1	ITEM 2	ITEM 3	ITEM 4
<b>Subject:1</b>				
True	.42	.27	.03	.36
1PL	.32	.30	.19	.32
2PL	.30	.30	.08	.30
3PL	.26	.25	.04	.23
<b>Subject:2</b>				
True	.32	.26	.05	.26
1PL	.26	.25	.15	.10
2PL	.27	.27	.05	.26
3PL	.26	.26	.05	.23
<b>Subject:3</b>				
True	.19	.28	.11	.20
1PL	.22	.21	.12	.07
2PL	.27	.27	.05	.26
2PL	.27	.28	.10	.24
<b>Subject:4</b>				
True	.50	.44	.47	.60
1PL	.55	.53	.38	.55
2PL	.46	.46	.50	.53
3PL	.46	.43	.46	.41
<b>Subject:5</b>				
True	.89	.77	.88	.95
1PL	.88	.88	.79	.88
2PL	.72	.72	.98	.84
3PL	.91	.73	.85	.96

TABLE 2. PARAMETERS OF OPERATIONAL MODELS  $\{\eta_0\}$ 

<u>1-PL</u>			
Item	b	a	c
1	.9	.7	0
2	1.0	.7	0
3	1.9	.7	0
4	.9	.7	0

<u>2-PL</u>			
1	1.4	.5	0
2	1.4	.5	0
3	1.1	1.7	0
4	.9	.7	0

<u>3-PL</u>			
1	1.3	3.2	.26
2	1.6	1.9	.25
3	1.1	2.1	.03
4	1.1	3.5	.23

TABLE 3. ASYMPTOTIC PARAMETERS OF MLE'S

Model	(1) $\theta_1$	(2) <sup>†</sup> $\theta_0$	(3) $-\mathbf{m}'(\theta_0, \eta_0)$	(4) <sup>*</sup> $E\mathbf{K}(\eta_0, \eta\hat{\theta}_0)$	(5) $\mathbf{K}(\eta_0, \eta\theta_0)$	(6) <sup>*</sup> $-\frac{1}{2}\mathbf{m}'(\theta_0, \eta_0)\mathbf{n}^{-1}\sigma_0^2$	(7) $\sigma_0^2$	(8) $I(\theta_0)^{-1}$
<b>Subject:1</b>								
1PL	-2	-2(3)	.10	.07	.04	.03	9.47	10.26
2PL	-2	-3(3)	.10	.04	.02	.02	7.61	9.76
3PL	-2	-1.2(1)	.00 <sup>+</sup>	.03	.03	.00 <sup>+</sup>	400.00	400.00
<b>Subject:2</b>								
1PL	-1	-.6(2)	.10	.07	.04	.03	8.68	10.26
2PL	-1	-.6(1.5)	.09	.03	.00 <sup>+</sup>	.03	11.42	11.76
3PL	-1	-.8(2)	.01	.04	.00 <sup>+</sup>	.04	133.33	133.33
<b>Subject:3</b>								
1PL	0	-.9(1)	.08	.06	.03	.03	13.33	13.33
2PL	0	-.6(1.5)	.09	.06	.02	.04	15.92	11.76
3PL	0	-.1(3)	.06	.04	.01	.03	18.06	16.67
<b>Subject:4</b>								
1PL	1	1.2(4)	.12	.04	.01	.03	8.16	8.16
2PL	1	1.1(4)	.24	.05	.02	.03	4.12	4.12
3PL	1	1.0(4)	.96	.05	.02	.03	1.03	1.04
<b>Subject:5</b>								
1PL	2	3.8(5)	.06	.06	.03	.03	14.08	16.00
2PL	2	3.3(5)	.06	.14	.07	.07	36.36	18.18
3PL	2	1.9(5)	.60	.03	.00 <sup>+</sup>	.03	1.68	1.66

\* n=16

† Rank order appears in parenthesis.

Department or Bureau	Civilian Agencies	Private Sector	Private Sector	Private Sector
12 Defense Technical Information Center Cameron Station, Bldg 5 Alexandria, VA 22304 Attn: IC	1 Dr. Patricia A. Butler HTE-004 Bldg, Stop 67 1200 19th St., NW Washington, DC 20208	1 Dr. Erling B. Andersen Department of Statistics Statensstatistisk 1435 Copenhagen DENMARK	1 Dr. Norman Cliff Dept. of Psychology Univ. of So. California University Park Los Angeles, CA 90007	1 Dr. Robert Blaser Learning Research & Development Center University of Pittsburgh 3939 O'Hara Street PITTSBURGH, PA 15260
1 Dr. Anita Lancaster Accession Policy DAB/MLB/MLP/MLP Perleap, Room 321 Washington, DC 20301	1 Dr. Vera B. Barry Personnel R&D Center Office of Personnel Management 1900 E Street NW Washington, DC 20415	1 Dr. Isaac Bejar Educational Testing Service Princeton, NJ 08506	1 Dr. Susan Emertson PSYCHOLOGY DEPARTMENT UNIVERSITY OF KANSAS Lawrence, KS 66045	1 Dr. Marvin D. Black 217 Stone Hall Cornell University Ithaca, NY 14853
1 Dr. Jerry Lohaus DAB/MLB Washington, DC 20301	1 Mr. Thomas A. Mara U. S. Coast Guard Institute P. O. Substation 18 Oklahoma City, OK 73169	1 Dr. Nemucha Birnbaum School of Education Tel Aviv University Tel Aviv, Ramat Aviv 60978 Israel	1 ERIC Facility-Acquisitions 4823 Rugby Avenue Bethesda, MD 20814	1 Dr. Bert Green Johns Hopkins University Department of Psychology Charles & 34th Street Baltimore, MD 21218
1 Dr. Clarence McCornick H&MPC/MLP 2500 Green Bay Road Naperville, IL 60564	1 Dr. Joseph L. Young, Director Henry & Cognitive Processes National Science Foundation Washington, DC 20550	1 Dr. Werner Birke Personalstatistik der Bundeswehr B-5000 Töln 90 WEST GERMANY	1 Dr. Benjamin A. Fairbank, Jr. Performance Metrics, Inc. 5825 Callaghan Suite 225 San Antonio, TX 78228	1 Dr. James E. Greene University of California, Berkeley Department of Education Berkeley, CA
1 Military Assistant for Training and Personnel Techniques Office of the Under Secretary of Defense for Research & Engineering Room 3B79, The Pentagon Washington, DC 20301		1 Dr. R. Berre, J. Bach Department of Education University of Chicago Chicago, IL 60637	1 Dr. Leonard Feldt Lindquist Center for Research University of Iowa Iowa City, IA 52242	1 Dipl. Pad. Michael M. Hahn Universität Bielefeld Erziehungswissenschaftliches Inst. II Bielefeld, Germany
1 Dr. W. Steve Seider Office of the Assistant Secretary on Defense R&E & 2B79, The Pentagon Washington, DC 20301		1 Dr. Arnold Bonner Section of Psychological Research Department of Psychology Casernes Petits Chateau CNR 1000 Brussels Belgium	1 Univ. Prof. Dr. Bernhard Fischer Liebigstrasse 27 A 1010 Vienna AUSTRIA	1 Dr. Ron Hambleton School of Education University of Massachusetts Amherst, MA 01002
1 Dr. Robert A. Bishop U.S. Army Institute for the Behavioral and Social Sciences 5001 Eisenhower Avenue Alexandria, VA 22304		1 Dr. Robert Brennan American College Testing Program P. O. Box 168 Iowa City, IA 52242	1 Professor Donald Fitzgerald University of New England Armidale, New South Wales 2351 AUSTRALIA	1 Prof. Lutz F. Hornig Universität Bielefeld Erziehungswissenschaftliches Inst. II Bielefeld, Germany
		1 Dr. Glenn Brvar 6208 Poe Road Bethesda, MD 20817	1 Dr. Dieter Fletcher University of Oregon Department of Computer Science Eugene, OR 97402	1 Dr. Paul Horst 677 6 Street, B184 Chula Vista, CA 92010
		1 Dr. Ernest B. Cadotte 307 Stoney University of Tennessee Knoxville, TN 37916	1 Dr. John R. Fredericksen Box: Berne & Hesser 50 Roulier Street Cambridge, MA 02128	1 Dr. Lloyd Humphreys Department of Psychology University of Illinois 602 East Daniel Street Champaign, IL 61820
		1 Dr. John B. Carroll 409 Elliott Rd. Chapel Hill, NC 27514	1 Dr. Janice Gifford University of Massachusetts School of Education Amherst, MA 01002	
			1 Dr. Stephen Duncan Lindquist Center for Measurement University of Iowa Iowa City, IA 52242	

# NAVY

1. Code 4711  
 1. Dr. Dick McShaw  
 WADP-125  
 Arlington Annex  
 Room 2024  
 Washington, DC 20380

1. Dr. Nick Bond  
 Office of Naval Research  
 Liaison Office, Far East  
 480 San Francisco, CA 94302

1. Dr. Alexander Berry  
 Applied Psychology  
 Measurement Division  
 NMD  
 NMD Persakia, S. 22508

1. Dr. Robert Braun  
 WADP-125  
 Code 4711  
 Dr. James F. 22508

1. Dr. Robert Carr  
 WADP-125  
 Washington, DC 20380

1. Dr. Stanley Collier  
 Office of Naval Research  
 800 M. Quincy St.  
 Arlington, VA 22217

1. Dr. Mike Corral  
 Office of Naval Research  
 800 M. Quincy St.  
 Arlington, VA 22217

1. Dr. John Ellis  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Dr. PAT FERBER  
 Code 4711  
 WADP-125  
 San Diego, CA 92152

1. Dr. Paul Foley  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Dr. Rebecca Heller  
 Navy Personnel R&D Center (Code 4711)  
 Building 4C  
 Brest Lakes NTC, IL 60088

# NAVY

1. Dr. Dick McShaw  
 WADP-125  
 Arlington Annex  
 Room 2024  
 Washington, DC 20380

1. Dr. Norman J. Kerr  
 Chief of Naval Education and Training  
 Code 400  
 Pensacola, FL 32508

1. Dr. Leonard Krueger  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Dr. David Lane  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Dr. William L. McChesney  
 Chief of Naval Education and Training  
 Naval Air Station  
 Pensacola, FL 32508

1. Dr. James McBride  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Dr. William Montague  
 WADP-125  
 San Diego, CA 92152

1. Dr. Kathleen Morris  
 Navy Personnel R&D Center (Code 4711)  
 San Diego, CA 92152

1. Library, Code 4711  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Technical Director  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Personnel & Training Research Program  
 Code 4829  
 Office of Naval Research  
 Arlington, VA 22217

# NAVY

1. Dr. Drew Sands  
 WADP-125  
 San Diego, CA 92152

1. Dr. Mary Schmitt  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Dr. Alfred F. Seale  
 Senior Scientist  
 Code 79  
 Naval Training Equipment Center  
 Orlando, FL 32812

1. Dr. Richard Sime  
 Liaison Scientist  
 Office of Naval Research  
 Branch Office, London  
 Box 39  
 FPO New York, NY 09506

1. Dr. Richard Sprensen  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Dr. Brad Swenson  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Dr. Frank Vicino  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Dr. Edward Weagar  
 Office of Naval Research (Code 4113)  
 800 North Quincy Street  
 Arlington, VA 22217

1. Dr. Ronald Weitzman  
 Naval Postgraduate School  
 Department of Administrative  
 Sciences  
 Monterey, CA 97940

1. Dr. Douglas Wetzel  
 Code 12  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Dr. MARTIN F. WISKOFF  
 NAVY PERSONNEL R&D CENTER  
 SAN DIEGO, CA 92152

# NAVY

1. Dr. John H. Wolfe  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Dr. Wallace Mulvey, III  
 Navy Personnel R&D Center  
 San Diego, CA 92152

1. Col. Ray Laidich  
 Headquarters, Marine Corps  
 WFO  
 Washington, DC 20380

1. Dr. William E. Nordbrock  
 FRC-ABCD Box 25  
 APO, NY 09710

1. Dr. Marcio F. O'Neil, Jr.  
 Director, Training Research Lab  
 Navy Research Institute  
 5001 Eisenhower Avenue  
 Alexandria, VA 22332

1. Commander, U.S. Army Research Institute  
 for the Behavioral & Social Sciences  
 ATTN: PERI-88 (Dr. Judith Orsano)  
 5001 Eisenhower Avenue  
 Alexandria, VA 22332

1. Major Frank Yohannan, USMC  
 Headquarters, Marine Corps  
 (Code 4711)  
 Washington, DC 20380

# NAVY

1. Dr. Kent Edson  
 Navy Research Institute  
 5001 Eisenhower Blvd.  
 Alexandria, VA 22332

1. Dr. Cressie Martin  
 Navy Research Institute  
 5001 Eisenhower Blvd.  
 Alexandria, VA 22332

1. Dr. Karen Mitchell  
 Navy Research Institute  
 5001 Eisenhower Blvd.  
 Alexandria, VA 22332

1. Dr. William E. Nordbrock  
 FRC-ABCD Box 25  
 APO, NY 09710

1. Dr. Marcio F. O'Neil, Jr.  
 Director, Training Research Lab  
 Navy Research Institute  
 5001 Eisenhower Avenue  
 Alexandria, VA 22332

1. Commander, U.S. Army Research Institute  
 for the Behavioral & Social Sciences  
 ATTN: PERI-88 (Dr. Judith Orsano)  
 5001 Eisenhower Avenue  
 Alexandria, VA 22332

1. Dr. Robert Ross  
 U.S. Army Research Institute for the  
 Social and Behavioral Sciences  
 5001 Eisenhower Avenue  
 Alexandria, VA 22332

1. Dr. Robert Sasser  
 U.S. Army Research Institute for the  
 Behavioral and Social Sciences  
 5001 Eisenhower Avenue  
 Alexandria, VA 22332

1. Dr. Joyce Shields  
 Navy Research Institute for the  
 Behavioral and Social Sciences  
 5001 Eisenhower Avenue  
 Alexandria, VA 22332

1. Dr. Hilda King  
 Navy Research Institute  
 5001 Eisenhower Ave.  
 Alexandria, VA 22332

# AIR FORCE

1. Dr. Kent Edson  
 Navy Research Institute  
 5001 Eisenhower Blvd.  
 Alexandria, VA 22332

1. Dr. Cressie Martin  
 Navy Research Institute  
 5001 Eisenhower Blvd.  
 Alexandria, VA 22332

1. Dr. Karen Mitchell  
 Navy Research Institute  
 5001 Eisenhower Blvd.  
 Alexandria, VA 22332

1. Dr. William E. Nordbrock  
 FRC-ABCD Box 25  
 APO, NY 09710

1. Dr. Marcio F. O'Neil, Jr.  
 Director, Training Research Lab  
 Navy Research Institute  
 5001 Eisenhower Avenue  
 Alexandria, VA 22332

1. Commander, U.S. Army Research Institute  
 for the Behavioral & Social Sciences  
 ATTN: PERI-88 (Dr. Judith Orsano)  
 5001 Eisenhower Avenue  
 Alexandria, VA 22332

1. Dr. Robert Ross  
 U.S. Army Research Institute for the  
 Social and Behavioral Sciences  
 5001 Eisenhower Avenue  
 Alexandria, VA 22332

1. Dr. Robert Sasser  
 U.S. Army Research Institute for the  
 Behavioral and Social Sciences  
 5001 Eisenhower Avenue  
 Alexandria, VA 22332

1. Dr. Joyce Shields  
 Navy Research Institute for the  
 Behavioral and Social Sciences  
 5001 Eisenhower Avenue  
 Alexandria, VA 22332

1. Dr. Hilda King  
 Navy Research Institute  
 5001 Eisenhower Ave.  
 Alexandria, VA 22332

Private Sector	Private Sector	Private Sector	Private Sector	Private Sector
1 Dr. Steven Hanks Department of Education University of Alberta Edmonton, Alberta Canada	1 Dr. Charles Lewis Facilitator Social Interchange Rijksuniversiteit Bruggen Oude Intergratiestraat 23 9712SC Bruggen Netherlands	1 Dr. Marvin R. Hewitt 356 Lindquist Center for Measurement University of Iowa Iowa City, IA 52242	1 Lowell Schuer Psychological & Quantitative Foundations College of Education University of Iowa Iowa City, IA 52242	1 Dr. Maurice Tatsuoka 220 Education Bldg 1310 S. Sixth St. Champaign, IL 61820
1 Dr. Jack Hunter 2122 College St. Lansing, MI 48906	1 Dr. Robert Litan College of Education University of Illinois Urbana, IL 61801	1 Dr. James Olson BICM, Inc. 1875 South State Street Brew, UT 84057	1 Dr. Kazuo Shigenasu 7-4-24 Tugunua-Kagan Fajutsua 251 JAPAN	1 Dr. David Thissen Department of Psychology University of Kansas Lawrence, KS 66044
1 Dr. Wayne Hahn College of Education University of South Carolina Columbia, SC 29208	1 Dr. Robert Lockman Center for Naval Analysis 200 North Beauregard St. Alexandria, VA 22311	1 Wayne R. Patience American Council on Education 883 Testing Service, Suite 20 One Dupont Circle, NW Washington, DC 20036	1 Dr. William Sias Center for Naval Analysis 200 North Beauregard Street Alexandria, VA 22311	1 Dr. Gary Thomason University of Illinois Department of Educational Psychology Champaign, IL 61820
1 Dr. Douglas H. Jones Advanced Statistical Technologies Corporation 10 Trafalgar Court Lawrenceville, NJ 08108	1 Dr. Frederic M. Lord Educational Testing Service Princeton, NJ 08541	1 Dr. James Paulson Dept. of Psychology Portland State University P.O. Box 751 Portland, OR 97207	1 Dr. M. Wallace Sinalko Program Director Rampower Research and Advisory Services Seafordman Institution 801 North Pitt Street Alexandria, VA 22314	1 Dr. Robert Tautakana Department of Statistics University of Missouri Columbia, MO 65201
1 Professor John A. Reats Department of Psychology The University of Newcastle N.S.W. 2308 AUSTRALIA	1 Dr. James Lussien Department of Psychology University of Western Australia Medlands W.A. 6009 AUSTRALIA	1 Dr. James A. Paulson Portland State University P.O. Box 751 Portland, OR 97207	1 Dr. Paul Spactman University of Missouri-Columbia Department of Statistics Columbia, MO 65201	1 Dr. Leifard Tucker University of Illinois Department of Psychology 403 E. Daniel Street Champaign, IL 61820
1 Dr. William Koch University of Texas-Austin Measurement and Evaluation Center Austin, TX 78702	1 Dr. Barry Marco Stop 31-E Educational Testing Service Princeton, NJ 08541	1 Dr. Mark D. Reclase ACT P. O. Box 168 Iowa City, IA 52242	1 Martha Stocking Educational Testing Service Princeton, NJ 08541	1 Dr. V. R. R. Uppluri Union Carbide Corporation Nuclear Division P. O. Box Y Oak Ridge, TN 37830
1 Dr. Thomas Leonard University of Wisconsin Department of Statistics 1210 West Barton Street Madison, WI 53705	1 Dr. Robert McKinley University of Toledo Dept of Educational Psychology Toledo, OH 43606	1 Dr. Lawrence Ruder 403 Elm Avenue Tahara Park, MD 20012	1 Dr. David Vale Assessment Systems Corporation 2233 University Avenue Suite 310 St. Paul, MN 55114	1 Dr. George Wong Biostatistics Laboratory Memorial Sloan-Kettering Cancer Center 1275 York Avenue New York, NY 10021
1 Dr. Alan Leopold Learning Rpt Center University of Pittsburgh 395 O'Hara Street Pittsburgh, PA 15260	1 Dr. Barbara Means Human Resources Research Organization 300 North Washington Alexandria, VA 22314	1 Dr. J. Ryan Department of Education University of South Carolina Columbia, SC 29208	1 Dr. Howard Weiner Division of Psychological Studies Educational Testing Service Princeton, NJ 08540	1 Dr. Mends Yen CTR/McGraw Hill Del Monte Research Park Monterey, CA 95940
1 Dr. Michael Levine Department of Educational Psychology 210 Education Bldg. University of Illinois Champaign, IL 61801	1 Dr. Robert Mislevy Educational Testing Service Princeton, NJ 08541	1 PPOE. EUMIKO SPEJINA DEPT. OF PSYCHOLOGY UNIVERSITY OF TENNESSEE KNOXVILLE, TN 37916	1 Dr. Ming-Mei Wang Lindquist Center for Measurement University of Iowa Iowa City, IA 52242	
	1 Dr. M. Alan Nicomander University of Oklahoma Department of Psychology Oklahoma City, OK 73069	1 Frank L. Schardt Department of Psychology Bldg. 66 George Washington University Washington, DC 20052	1 Dr. Brian Waters HueRHO 300 North Washington Alexandria, VA 22314	



**END**

**FILMED**

3-86

**DTIC**